

COXI-STAT™ COCCIDIOCIDAL SOLUTION FOR POULTRY

ACTIVE CONSTITUENT: 400g/L AMPROLIUM (as Amprolium Hydrochloride)

Easy, effective coccidiosis protection in every drop

For the prevention and treatment of coccidiosis in broilers, broiler breeders, replacement pullets, turkeys, ducks and pigeons.

- Comprehensive Protection – Prevents and treats coccidiosis
- Broad-Spectrum Control – Targets major *Eimeria* species, including *E. tenella*, *E. necatrix*, and *E. acervulina*
- Easy Administration – Just mix in drinking water!
- Suitable for multiple poultry types



COXI-STAT™	
Active Constituent	400g/L AMPROLIUM (as Amprolium Hydrochloride)
Species	Poultry
Route of Administration	Oral, Drinking water
Dose rate - Prevention	15mL/25L of drinking water for 24 hours every 4th day
Dose rate - Treatment	15mL/25L of water for 5-7 days, reduce to 7.5mL/25L for a further 5-7 days, then to 4mL/25L for 5-7 days
WHP	Zero Days
ESI	Not established

COXI-STAT™ is a trademark of Abbey Laboratories Pty Ltd

COXI-STAT™ COCCIDIOCIDAL SOLUTION FOR POULTRY

ACTIVE CONSTITUENT: 400g/L AMPROLIUM (as Amprolium Hydrochloride)



For the prevention and treatment of coccidiosis in broilers, broiler breeders, replacement pullets, turkeys, ducks and pigeons.

INDICATIONS

Amprolium belongs to the class of methylpyridines and is a thiamine antagonist. It is used to prevent and treat intestinal coccidiosis by blocking the thiamine transporter (Competitive inhibition of thiamine uptake).

COXI-STAT™ Coccidiocidal Solution for Poultry is administered through the drinking water as an aid in the prevention and treatment of economically important strains of coccidiosis in poultry including *E. tenella*, *E. necatrix* and most strains of *E. acervulina*. Resistance may develop to any chemical.

DIRECTIONS FOR USE

RESTRAINTS

For professional use only (veterinary surgeons and assistants, and farmers).

DOSAGE AND ADMINISTRATION

Discard unused portion of the product within four months of first broaching. Replace medicated drinking water after 24 hours.

DOSAGE FOR PREVENTION: Add 15mL per 25L of drinking water (240mg amprolium per litre) for 24 hours every fourth day during periods of likely infection. This action helps control the coccidia by interrupting the life cycle and encourages immunity to develop.

DOSAGE FOR TREATMENT: Add 15mL per 25L of drinking water (240mg amprolium per litre) CONTINUOUSLY for 5 to 7 days, reduce this to 7.5mL per 25L for a further 5 to 7 days, then to 4mL per 25L for another 5 to 7 days. This gradual reduction in the level of treatment assists the birds to develop immunity while maintaining some level of protection.

ADMINISTRATION: Sufficient access to the water supply system should be available for the birds to be treated to ensure adequate water consumption. No other source of drinking water should be available during the medication period.

WITHHOLDING PERIODS

Zero (0) days.

SAFETY DIRECTIONS

Will damage eyes and skin. The product is strongly acidic. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. When opening the container and using the product wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow-length chemical resistant gloves and face shield or goggles. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

First aid is not generally required. If in doubt, contact a Poisons Information Centre. Phone Australia 13 11 26.

DISPOSAL

For 1L: Dispose of container by wrapping with paper and putting in garbage.

For 5L: Triple-rinse container into the medicated water. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean container to recycler or designated collection point. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

STORAGE

Store below 25°C (air conditioning). Do not freeze. Protect from light. KEEP OUT OF REACH OF CHILDREN.

APVMA Approval Number: 94310/146760